HEALTHEAST MEDICAL TRANSPORTATION MEDICAL OPERATIONS MANUAL

LIDOCAINE

Drug Classification: Antiarrhythmic, local anesthetic

BACKGROUND

Pharmacodynamics

Blocks sodium channels, increasing the recovery period after repolarization; suppresses automaticity in the His-Purkinje system and depolarization in the ventricles.

Indications

- Pain with medication or fluid administration through IO.
- V-fib or V-tach arrest, if allergic to or currently taking amiodarone (Cordarone).
- Pre-treatment for suspected increased intracranial pressure prior to DFAM.

Contraindications

- Allergy or known hypersensitivity to the medication.
- Severe degrees of SA, AV, or intraventricular block.

Cautions

• May cause injection site pain and/or nausea and vomiting.

DOSAGE and ADMINISTRATION

ADULT	PEDIATRICS (less than 60 kg)
For pain management with IO	For pain management with IO
Administer 50 mg slowly IO over 2 minutes followed by a fluid flush.	 Administer 0.5 mg/kg slowly IO over 2 minutes followed by a fluid flush.
For V-Fib or V-Tach without a pulse	For V-Fib or V-Tach without a pulse
1. Administer 1 mg/kg IV/IO.	1. Administer 1 mg/kg IV/IO.
2. Administer additional doses at 0.5 mg/kg every 5 min up to a total dose of 3 mg/kg.	Administer additional doses at 0.5 mg/kg every 5 min up to a total dose of 3 mg/kg.
For DFAM pre-treatment in suspected increased cranial pressure	For DFAM pre-treatment in suspected increased cranial pressure
1. 1 mg/kg IV/IO max dose 100 mg for suspected increased intracranial pressure.	1 mg/kg IV/IO max dose 100 mg for suspected increased intracranial pressure.

ADVERSE REACTIONS/SIDE EFFECTS

Seizures, cardiac arrhythmias, hypotension

Return to TOC Lidocaine: Page 1 of 2 Version 2016.1

HEALTHEAST MEDICAL TRANSPORTATION MEDICAL OPERATIONS MANUAL

NOTES

REFERENCE GUIDELINE/PROCEDURE:

2K Ventricular Ectopy 7F1 DFAM 7R Intraosseous (IO) Access

Return to TOC Lidocaine: Page 2 of 2 Version 2016.1